arXiv:1201.5060v2 [quant-ph] 6 Feb 2012

Storing flux qubits in the quantum RAM of binary Bose-Einstein condensates

Kelly R. Patton and Uwe R. Fischer
Seoul National University, Department of Physics and Astronomy
Center for Theoretical Physics, 151-747 Seoul, Korea

We consider the hybrid quantum system of an rf-SQUID interacting via its magnetic field with a two-component Bose-Einstein condensate. It is shown that the effective coupling between the quantized SQUID flux and the atomic hyperfine states scales with the total number of particles in the condensate. We demonstrate that this renders the two-component condensate a prospective architecture for rapidly accessible storage and retrieval of the quantum information encoded in the flux qubits with high transfer fidelity.

PACS numbers: 03.67.-a, 03.75.Mn, 85.25.Dq

The concept of hybrid quantum systems, combining various advantages (and compensating for deficiencies) of either subsystem is rapidly gaining momentum as a concrete tool for the manipulation of quantum information [1]. Various examples for such hybrid systems include cavity-QED with Bose-Einstein condensates (BECs) [2], nanomechanical resonators coupled to superconductors [3, 4], and ultracold atoms coupled to a superconducting waveguide cavity [5].

Using the hybrid system as a quantum computer, one part acts as the quantum central processing unit (CPU), while the other component is used as the quantum analog of a random access memory (RAM). A fundamental application question is, then, how to quickly store quantum information in the RAM which has to possess a low decoherence rate, and equally rapidly retrieve it to make quantum computations in the CPU. In the present study, we investigate the hybrid system shown in Fig. 1, and show that it is especially suitable for the particular application just described. An rf-SQUID is coupled magnetically to a two-component miscible BEC, where two hyperfine components, equally populated with $N/2$ atoms, interpenetrate. Similar hybrids, integrating cooling techniques for both ultracold atoms as well as superconductors, have already been demonstrated to be experimentally feasible [6–8].

While the collective Rabi exchange process induced by flux qubit operations with the SQUID pseudospin [9] flips only a single atom pseudospin in the BEC at a time, the SQUID–BEC coupling is enhanced by a factor $\sqrt{N/2}$ contributed by *each* hyperfine component (N is total number of atoms in the BEC). This is due to the bosons in each hyperfine level being indistinguishable: We do not know *which* of the $N/2$ single spins has been flipped of either component of the BEC. The indistinguishability of the atoms in the condensate therefore creates an $\mathcal{O}(N)$ speedup of the flux qubit transfer process. In addition, we will take advantage of the fact that SQUID and BEC can rapidly be brought into and out of resonance by tuning the flux qubit tunneling frequency on very short time scales, on the order of nanoseconds [10, 11]. As a result, we shall demonstrate that several complete transfer cycles of the SQUID qubit state are possible within the decoherence time of the SQUID. The advantages of this quantum RAM [12] over other (solid-state based) proposals for qubit storage in hybrid systems [13–18], are that the quantum information encoded in the hyperfine qubit lives for a long time before it decoheres without, e.g., the need for further transformation to nuclear spin states, and that it can rapidly be transferred back and forth between SQUID and BEC. Finally, the quantum RAM used in the following has decoherence time scales exceeding by several orders of magnitude those currently achievable for the parent flux qubits [19].

We first discuss the theoretical description of the two subsystems and then derive an effective Hamiltonian, taking into account their coupling. The basic assumption underlying the existence of flux qubits is that the flux through a superconducting ring is a quantum variable governed by an effective potential with two minima corresponding to clock- and counterclockwise circulating electrons. In the low-energy limit, in which the two-mode


Two-component BEC
Ω
Hyperfine qubit
$\alpha|0\rangle + \beta|1\rangle$
$\boldsymbol{B}(\boldsymbol{r},t)$
State
Swap
Flux qubit
$\alpha|\mathrm{L}\rangle + \beta|\mathrm{R}\rangle$
SQUID

FIG. 1. A flux qubit created from superposing left- and right-going current states of a rf-SQUID is coupled by its magnetic field to the hyperfine transition of a two-component BEC, where the Rabi process periodically flips one atom from a given hyperfine spin state to the other. The representation is not to scale.

approximation applies, the SQUID Hamiltonian is [20]

$$\hat{H}_{\mathrm{SQUID}} = \frac{\varepsilon}{2}\sigma_z - \frac{\Delta}{2}\sigma_x, \tag{1}$$

where ε is the energy difference (the effective flux potential offset) between the eigenstates $|\mathrm{L}\rangle$ and $|\mathrm{R}\rangle$ in the left- and right-going current basis, and Δ represents the tunneling rate through the potential barrier. In the left-right current eigenbasis, a *current operator* can be defined as follows, $\hat{I} \simeq I\sigma_z$, where $I = \langle \mathrm{L}|\hat{I}(\hat{\Phi})|\mathrm{L}\rangle \simeq -\langle \mathrm{R}|\hat{I}(\hat{\Phi})|\mathrm{R}\rangle$ (exact equality holding when $\epsilon = 0$). The current operator is related to the flux operator through the SQUID loop by $\hat{I} = \frac{\hat{\Phi}-\Phi_{\mathrm{ex}}}{L}$ with Φ_{ex} the classical externally applied flux and L the loop inductance [21]. The magnetic vector potential of a loop of radius R in the dipole approximation and in polar coordinates reads $\hat{\boldsymbol{A}} \simeq -\frac{\mu_0 R^2 \hat{I}}{4r^2}\sin\theta \boldsymbol{e}_\phi$. Thus the qubit magnetic field is given by $\hat{\boldsymbol{B}} = \nabla \times \hat{\boldsymbol{A}} = \boldsymbol{B}(\boldsymbol{r})\sigma_z$.

The interacting Hamiltonian for a trapped two-component BEC with field operator components $\hat{\Psi}_\sigma$, where the pseudospin $\sigma = (\downarrow, \uparrow)$ labels the hyperfine index, takes the form ($\hbar = 1$)

$$\begin{aligned}\hat{H}_{\mathrm{BEC}} = &\sum_{\sigma=\uparrow,\downarrow} \int \mathrm{d}^D r\, \hat{\Psi}^\dagger_\sigma(\boldsymbol{r}) \left[-\frac{\nabla^2}{2m} + \omega_\sigma + V(\boldsymbol{r})\right] \hat{\Psi}_\sigma(\boldsymbol{r}) \\ &+ \frac{1}{2}\sum_{\sigma,\sigma'} \mathrm{g}_{\sigma,\sigma'} \int \mathrm{d}^D r\, \hat{\Psi}^\dagger_\sigma(\boldsymbol{r}) \hat{\Psi}^\dagger_{\sigma'}(\boldsymbol{r}) \hat{\Psi}_{\sigma'}(\boldsymbol{r}) \hat{\Psi}_\sigma(\boldsymbol{r}),\end{aligned} \tag{2}$$

where $V(\boldsymbol{r})$ is the trapping potential, assumed to be independent of spin, ω_σ corresponds to the hyperfine energy levels, and $\mathrm{g}_{\sigma,\sigma'}$ are interaction couplings.

The magnetic coupling of the BEC to the SQUID is given by the canonical expression

$$\hat{H}_{\mathrm{int}} = -\sum_{\sigma,\sigma'} \int \mathrm{d}^D r\, \hat{\Psi}^\dagger_\sigma(\boldsymbol{r}) \left[\boldsymbol{\mu}\right]_{\sigma,\sigma'} \hat{\Psi}^\dagger_{\sigma'}(\boldsymbol{r}) \otimes \hat{\boldsymbol{B}}(\boldsymbol{r}), \tag{3}$$

multiplying the magnetic moment density of atoms with the local magnetic field, and integrating over space. Here, $\boldsymbol{\mu}$ is the total magnetic moment matrix of atoms in the σ basis. The total Hamiltonian is then

$$\hat{H} = \hat{H}_{\mathrm{BEC}} \otimes \mathbb{1} + \mathbb{1} \otimes \hat{H}_{\mathrm{SQUID}} + \hat{H}_{\mathrm{int}}, \tag{4}$$

where $\mathbb{1}$ is the identity matrix in either BEC or SQUID space (corresponding to the first and second vector product entry, respectively).

We now derive an approximation to the full Hamiltonian, which represents the coupling of the flux qubit to a large two-component BEC on the mean-field level. To this end, we first truncate the field operators using the standard mean-field approximation to just one mode for each species σ,

$$\hat{\Psi}_\sigma(\boldsymbol{r}) \simeq \phi_\sigma(\boldsymbol{r})\hat{a}_\sigma, \tag{5}$$

where $\phi_\sigma(\boldsymbol{r})$ are mean-field solutions for the interacting BEC, leading to coupled Gross-Pitaevskiĭ equations.

As we will see below, even with bosonic enhancement, the SQUID-BEC coupling is small compared to the hyperfine energy difference, so that their ratio is a small parameter. In terms of this parameter, the two systems are still weakly coupled. Along with the form of the interaction (3), which only couples states in the BEC Hilbert space that differ by a single pseudospin, this leads us to conclude that the full Hilbert space of the BEC can be approximated by a two-dimensional subspace; the "ground state" with $N_\downarrow$ and $N_\uparrow$ atoms and an "excited state" that differs by only a single atom of each spin. We therefore restrict ourselves to the following two BEC basis states,

$$\begin{aligned}|0\rangle_{\mathrm{B}} =& \frac{1}{\sqrt{N_\downarrow!}}\frac{1}{\sqrt{N_\uparrow!}}\left(\hat{a}^\dagger_\downarrow\right)^{N_\downarrow}\left(\hat{a}^\dagger_\uparrow\right)^{N_\uparrow}|\mathrm{vac}\rangle \\ |1\rangle_{\mathrm{B}} =& \frac{1}{\sqrt{(N_\downarrow-1)!}}\frac{1}{\sqrt{(N_\uparrow+1)!}}\left(\hat{a}^\dagger_\downarrow\right)^{N_\downarrow-1}\left(\hat{a}^\dagger_\uparrow\right)^{N_\uparrow+1}|\mathrm{vac}\rangle.\end{aligned} \tag{6}$$

In the weak coupling limit (in the sense described above), the dominant process is the periodic transfer of a single quantum of energy from one subsystem to the other, i.e., the Rabi cycle. The whole binary BEC appears in this limit, and within mean-field Eq. (5), as a giant two-level atom coupled to one "cavity" mode produced by the SQUID. We have, in addition, verified explicitly that processes flipping more than one spin at a time have negligibly small probability.

As the BEC Hamiltonian (2) conserves spin, its off-diagonal elements in the basis of (6) vanish, while the diagonal ones give the mean-field energies $\mathcal{E}_{N_\uparrow,N_\downarrow}, \mathcal{E}_{N_\uparrow+1,N_\downarrow-1}$. Using that $\mathcal{E}_{N_\uparrow+1,N_\downarrow-1} - \mathcal{E}_{N_\uparrow,N_\downarrow} \simeq \omega_\uparrow - \omega_\downarrow$, (the interaction renormalization of the hyperfine splitting being small compared to the level spacing), the interacting Hamiltonian (4), can finally be written as, omitting $\mathcal{O}(1/N)$ and constant terms,

$$\begin{aligned}\hat{H} \simeq& \frac{\omega_\uparrow - \omega_\downarrow}{2}\sigma_z \otimes \mathbb{1} + \mathbb{1} \otimes \left(\frac{\varepsilon'}{2}\sigma_z - \frac{\Delta}{2}\sigma_x\right) \\ &- \sqrt{N_\downarrow N_\uparrow}\begin{pmatrix} 0 & \boldsymbol{g}_{\uparrow,\downarrow}\cdot\boldsymbol{\mu}_{\uparrow,\downarrow} \\ \boldsymbol{g}_{\downarrow,\uparrow}\cdot\boldsymbol{\mu}_{\downarrow,\uparrow} & 0 \end{pmatrix} \otimes \sigma_z.\end{aligned} \tag{7}$$

The SQUID–BEC coupling matrix is expressed as

$$\boldsymbol{g}_{\sigma,\sigma'} = \int \mathrm{d}^D r\, \phi^*_\sigma(\boldsymbol{r})\boldsymbol{B}(\boldsymbol{r})\phi_{\sigma'}(\boldsymbol{r}), \tag{8}$$

and $\varepsilon' = \varepsilon + N_\uparrow \boldsymbol{g}_{\uparrow,\uparrow}\cdot\boldsymbol{\mu}_{\uparrow,\uparrow} + N_\downarrow \boldsymbol{g}_{\downarrow,\downarrow}\cdot\boldsymbol{\mu}_{\downarrow,\downarrow}$ is a renormalized level splitting of the SQUID.

A first major result of the present investigation is the $\mathcal{O}(N)$ enhancement of the effective coupling in the last term of Eq. (7). Using this effective Hamiltonian, it is easily shown that one can prepare the flux qubit in an arbitrary state and then pass it to the BEC; because of Rabi oscillations this state will then get passed back to the SQUID, with the process repeatable for many cycles due to the short duration of transfer. We first focus on controlledly passing the state to the BEC for it to remain there. This can be achieved by bringing the SQUID into and out of resonance with the given hyperfine splitting of the atoms. Here, we will crucially take advantage of the fact that one can dynamically change the flux tunneling barrier of the SQUID potential on a essentially arbitrarily rapid timescale within a large frequency window (Ref. [10] quotes a change of several GHz for Δ on timescales of nanoseconds). Changing the tunneling rate amounts to replacing $\Delta \to \Delta(t)$. The time dependence should be chosen such that the SQUID and the BEC are only in resonance, i.e., $\Delta(t) = \omega_\uparrow - \omega_\downarrow$ for exactly half a Rabi period when one transfers the flux qubit to the BEC. Outside of this window they should be far off resonance. One choice for this is simply

$$\Delta(t) = \Delta_0 W(t), \tag{9}$$

where $\Delta_0 = \omega_\uparrow - \omega_\downarrow$ and $W(t)$ ramps from an arbitrarily chosen off resonance value to 1 for half a Rabi period and then ramps back to its off resonance value. To remain in the qubit subspace, i.e., to not excite higher SQUID flux potential levels, the ramping time should remain slow compared to Δ_0^{-1}.

Assuming the SQUID is at the degeneracy point for simplicity, $\varepsilon' = 0$, the energy eigenstates of an isolated SQUID are $|0\rangle_\mathrm{S} = \frac{1}{\sqrt{2}}\big[|\mathrm{L}\rangle + |\mathrm{R}\rangle\big]$ and $|1\rangle_\mathrm{S} = \frac{1}{\sqrt{2}}\big[|\mathrm{L}\rangle - |\mathrm{R}\rangle\big]$. The computation basis for the combined system is then taken to be $|00\rangle = |0\rangle_\mathrm{B} \otimes |0\rangle_\mathrm{S}$, $|01\rangle = |0\rangle_\mathrm{B} \otimes |1\rangle_\mathrm{S}$, $|10\rangle = |1\rangle_\mathrm{B} \otimes |0\rangle_\mathrm{S}$, and $|11\rangle = |1\rangle_\mathrm{B} \otimes |1\rangle_\mathrm{S}$. For a system with equal intial populations, $N_\uparrow = N_\downarrow = N/2$, the Hamiltonian in the computation basis reads, setting $\Delta_0 = \omega_\uparrow - \omega_\downarrow$,

$$\hat{H}(t) \simeq \frac{\Delta_0}{2}\sigma_z \otimes \mathbb{1} + \mathbb{1} \otimes \frac{\Delta_0 W(t)}{2}\sigma_z - \begin{pmatrix} 0 & \Omega \\ \Omega^* & 0 \end{pmatrix} \otimes \sigma_x, \tag{10}$$

where $\Omega = \frac{N}{2}\boldsymbol{g} \cdot \boldsymbol{\mu}_{\uparrow,\downarrow}$ is enhanced by a factor $N/2$ over the single-particle coupling.

The Hamiltonian (10) is easily transformed into a cavity-QED type Hamiltonian. Here the two-component BEC corresponds to a two-level atom, while the SQUID is the cavity photon. For the SQUID we let $\frac{\Delta}{2}\sigma_z \to \Delta \hat{a}^\dagger \hat{a}$, where $\hat{a}, \hat{a}^\dagger$ are photon annihilation and creation operators. Assuming a real Rabi frequency Ω, we replace $\sigma_x \to \hat{a}^\dagger + \hat{a}$ for the SQUID and $\sigma_x \to \sigma_+ + \sigma_-$ for the BEC operators in the coupling term. We then have in rotating wave approximation [22]

$$\hat{H} = \frac{\Delta_0}{2}\sigma_z \otimes \mathbb{1} + \mathbb{1} \otimes \Delta \hat{a}^\dagger \hat{a} - \Omega\big(\sigma_- \otimes \hat{a}^\dagger + \sigma_+ \otimes \hat{a}\big), \tag{11}$$

which is a Jaynes-Cummings Hamiltonian cf., e.g., [23]. For the calculations presented below, we have verified that $\langle \hat{a}^\dagger \hat{a}\rangle \leq 1$ at any instant, so that at most one photon is in the SQUID mode.

We take as a concrete example a ^{87}Rb BEC of $N = 10^6$ atoms, which has been shown to give miscible two-component condensates when the states $|\downarrow\rangle = \big|5^2S_{1/2}, F = 1, m_F = -1\big\rangle$ and $|\uparrow\rangle = \big|5^2S_{1/2}, F = 2, m_F = -2\big\rangle$ are used [24]. For the purpose of our computation examples below, we set $\Delta_0 = 6.835\,\mathrm{GHz}$ equal to the hyperfine splitting without Zeeman shift between $|\downarrow\rangle$ and $|\uparrow\rangle$ (a non-degenerate two-level hyperfine qubit can be formed from these states by applying a small external magnetic field to lift the m_F degeneracy). The single-particle coupling strength is taken to be $|\boldsymbol{g} \cdot \boldsymbol{\mu}_{\uparrow,\downarrow}| \approx 20\,\mathrm{Hz}$, corresponding to a SQUID radius of $1\,\mu\mathrm{m}$, with a current of $1\,\mu\mathrm{A}$, and a BEC-SQUID distance of about $20\,\mu\mathrm{m}$. The fact that the latter distance is much larger than the SQUID radius implies that the detrimental effect of Meissner screening on the trapping is negligible [8], representing a major advantage over superconducting cavities, where the cold atom cloud is situated close to a large waveguide geometry [5]. The spin-flip lifetime due to current fluctuations should not be a limiting factor at distances of this order as well [25]; finally, the SQUID needs to be cooled to temperatures below 100 mK, to avoid disturbance of the qubit generation and transfer process by thermal photon emission.

An arbitrary state in the computation basis can be decomposed as the superposition

$$|\Psi\rangle = c_1|11\rangle + c_2|01\rangle + c_3|10\rangle + c_4|00\rangle. \tag{12}$$

The coefficients c_i satisfy the coupled Schrödinger equations $\partial_t c_i(t) = -i\hat{H}(t)c_i(t)$, for a given $c_i(t=0)$ initial condition. We now provide two examples for a transfer protocol, where the results are calculated numerically



FIG. 2. Top panel: Probabilities for finding the system in the state $|01\rangle$ and $|10\rangle$ as a function of time for two different ramping times. Left: Ramping time 5 ns with final (time-averaged) fidelity $F = 99.3\%$. Right: 10 ns ramping with final $F = 95.9\%$ (we fix $\Delta_0^{-1} = 0.15\mathrm{ns}$). Bottom panel: Final time-averaged fidelity as a function of ramp time.



FIG. 3. The top panel shows the transfer of a superposition state of the flux qubit $|\psi(t=0)\rangle = |0\rangle_{\rm B} \otimes 2^{-1/2}(|0\rangle_{\rm S} + |1\rangle_{\rm S}) = 2^{-1/2}(|00\rangle + |01\rangle)$ to the hyperfine qubit state $|\psi(t_{\rm f})\rangle = 2^{-1/2}(|0\rangle_{\rm B} + |1\rangle_{\rm B}) \otimes |0\rangle_{\rm S} = 2^{-1/2}(|00\rangle + |10\rangle)$. The middle panel shows the production of the state $|\psi\rangle = 2^{-1/2}(|01\rangle + |10\rangle)$ from an initial state $|01\rangle$. Finally, the bottom panel demonstrates a memory transfer process with a final fidelity of 99%. The ramping times for all figures are 5 ns.

without using the rotating wave approximation.

We start with a simple transfer protocol. If at time $t = 0$ the combined system is in the state $|\Psi\rangle = |01\rangle$, then $c_i(t=0) = 0$ for $i \neq 2$ and $c_2(t=0) = 1$. The probability that at some later time the system is found in the state $|10\rangle$ is $|c_3(t)|^2$, see Fig. 2, where we give examples of the fidelity $F = |\langle\Psi(\text{target})|\Psi\rangle|$, evaluating for this protocol to $F(t) = \left|a^*_{00}(t)c_{00}(t) + a^*_{10}(t)c_{10}(t)\right|$ writing the time evolution of the initial state as $|\Psi\rangle = \sum_{m,n} c_{m,n}(t)|mn\rangle$ and similarly that of the target state as $|\Psi(\text{target})\rangle = \sum_{m,n} a_{m,n}(t)|mn\rangle$. We define *ramp time* as the time interval it takes to raise Δ from 1 % above its value at $t \to -\infty$ to within 1 % of being on resonance, using a smooth tanh profile for $W(t)$ [26]. The bottom panel of Fig. 2 shows that F decreases with the ramp time thus defined. Rapid transfer (while staying within the limits of the two-mode approximation) therefore significantly improves on the fidelity.

More elaborate examples for flux qubit storage and retrieval are displayed in Fig. 3, where we show in the top panel that an almost perfect transfer (fidelity $F = 99.3\%$) of a flux qubit to the hyperfine qubit is possible within a short time (we have verified that an initial superposition state with phase difference $\pi/2$, $|\psi(t=0)\rangle = |0\rangle_{\rm B} \otimes 2^{-1/2}(|0\rangle_{\rm S} + i|1\rangle_{\rm S}) = 2^{-1/2}(|00\rangle + i|01\rangle)$ yields the same result). The middle panel of Fig. 3 shows that one can entangle the two qubits, i.e., produce a final state $|\psi\rangle = 2^{-1/2}(|01\rangle + |10\rangle)$, by preparing the system in the state $|01\rangle$ at $t = 0$, and then bring them into resonance for a quarter of a Rabi cycle. The latter process is in the example shown achieved with a fidelity of 98.8 %. Finally, the bottom panel demonstrates the quantum RAM in storage and retrieval action, by transferring first the initial qubit state to the BEC and then back to the SQUID by again tuning into resonance, showing that this concrete RAM-type process can be achieved as well with large fidelity, here $F = 99\%$.

In conclusion, we have demonstrated that rapid and efficient flux qubit storage and retrieval are feasible when an rf-SQUID is magnetically coupled to the hyperfine levels of a two-component BEC. We expect a periodic extension of the present hybrid to be conceivable, with atomic ensembles in a large-filling optical lattice located on top of an array of SQUIDs. With such a setup, the flux qubits can be entangled with each other via the condensates in the lattice sites.

Transferring back and forth flux qubits to long-lived hyperfine qubit states of atoms in a two-component BEC, for which one has rather precise control over the microscopic quantum physics, illuminates further whether the postulated quantum variable flux has fundamental character. This is directly related to long-standing intriguing questions on the existence of macroscopic quantum variables and thus the quantum–classical transition [27, 28]. Among future research directions, then, is the potential influence of microscopic variables of the SQUID on the qubit transfer process.

This research was supported by the NRF of Korea, grant Nos. 2010-0013103 and 2011-0029541. URF thanks D. Cano, J. Fortágh, R. Kleiner, and C. Zimmermann for helpful discussions.

[1] M. Wallquist, K. Hammerer, P. Rabl, M. Lukin, and P. Zoller, Phys. Scr. **T137** (2009) 014001.
[2] F. Brennecke *et al.*, Nature **450**, 268 (2007); K. Baumann, C. Guerlin, F. Brennecke, and T. Esslinger, Nature **464**, 1301 (2010).
[3] A. D. O'Connell *et al.*, Nature **464**, 697 (2010).
[4] A. N. Cleland and M. R. Geller, Phys. Rev. Lett. **93**, 070501 (2004).
[5] J. Verdú, H. Zoubi, Ch. Koller, J. Majer, H. Ritsch, and J. Schmiedmayer, Phys. Rev. Lett. **103**, 043603 (2009).
[6] T. Nirrengarten *et al.*, Phys. Rev. Lett. **97**, 200405 (2006); C. Roux *et al.*, Europhys. Lett. **81**, 56004 (2008).
[7] T. Mukai *et al.*, Phys. Rev. Lett. **98**, 260407 (2007).
[8] D. Cano, B. Kasch, H. Hattermann, R. Kleiner, C. Zimmermann, D. Koelle, and J. Fortágh, Phys. Rev. Lett. **101**, 183006 (2008); D. Cano *et al.*, Eur. Phys. J. D **63**, 17 (2011).
[9] I. Chiorescu, Y. Nakamura, C. J. P. M. Harmans, and J. E. Mooij, Science **299**, 1869 (2003).

[10] F. G. Paauw, A. Fedorov, C. J. P. M. Harmans, and J. E. Mooij, Phys. Rev. Lett. **102**, 090501 (2009).

[11] M. G. Castellano, F. Chiarello, P. Carelli, C Cosmelli, F. Mattioli, and G. Torrioli, New J. Phys. **12**, 043047 (2010).

[12] M. Blencowe, Nature **468**, 44 (2010).

[13] I. Chiorescu, N. Groll, S. Bertaina, T. Mori, and S. Miyashita, Phys. Rev. B **82**, 024413 (2010).

[14] D. I. Schuster *et al.*, Phys. Rev. Lett. **105**, 140501 (2010).

[15] Y. Kubo *et al.*, Phys. Rev. Lett. **105**, 140502 (2010); Phys. Rev. Lett. **107**, 220501 (2011).

[16] H. Wu *et al.*, Phys. Rev. Lett. **105**, 140503 (2010).

[17] X. Zhu *et al.*, Nature **478**, 221 (2011).

[18] M. Mariantoni *et al.*, Science **334**, 61 (2011).

[19] J. Bylander *et al.*, Nature Physics **7**, 565 (2011).

[20] Y. Makhlin, G. Schön, and A. Shnirman, Rev. Mod. Phys. **73**, 357 (2001).

[21] We note that operator notation for SQUID magnetic field variables exclusively refers to the flux qubit space, and there is no field quantization implied.

[22] Without the rotating wave approximation, we obtain the (more fundamental) Rabi model; I. I. Rabi, Phys. Rev. **49**, 324 (1936); **51**, 652 (1937).

[23] H. Walther, B. T. H. Varcoe, B.-G. Englert, and T. Becker, Rep. Prog. Phys. **69**, 1325 (2006).

[24] M. A. Hoefer, J. J. Chang, C. Hamner, and P. Engels, Phys. Rev. A **84**, 041605(R) (2011).

[25] R. Fermani, T. Müller, B. Zhang, M. J. Lim, and R. Dumke, J. Phys. B: At. Mol. Opt. Phys. **43**, 095002 (2010); B. Kasch *et al.*, New J. Phys. **12**, 065024 (2010).

[26] Explicitly, $W(t) = \Theta(k,t)\Theta(k,\tau - t) + \Delta_{\rm or}\Theta(k,-t) + \Delta_{\rm or}\Theta(k,t-\tau)$, where $\Theta(k,t) = \frac{1}{2} + \frac{1}{2}\tanh[kt]$, τ is the time the two systems are on resonance (typically for half a Rabi cycle), and $\Delta_{\rm or}$ is the off resonance value of the SQUID tunneling in units of Δ_0 (setting $\Delta_{\rm or} = \frac{1}{2}$ for all our calculations). The function $\Theta(k,t)$ is a smoothed step function, such that $\lim_{k\to\infty}\Theta(k,t) = \Theta(t)$; different ramp times are obtained by varying k.

[27] A. J. Leggett, J. Phys.: Condens. Matter **14**, R415 (2002).

[28] A. Fedorov, P. Macha, A. K. Feofanov, C. J. P. M. Harmans, and J. E. Mooij, Phys. Rev. Lett. **106**, 170404 (2011).